



International Baccalaureate®
Baccalauréat International
Bachillerato Internacional

Diploma Programme

Mathematics: analysis and approaches HL formula booklet

For use during the course and in the examinations
First examinations 2021

Version 1.0

Two decorative blue curves are present on the page. One is a thick, dark blue curve that starts on the left, rises to a peak, and then descends towards the right. The other is a thin, light blue curve that starts high on the left, descends to a minimum, and then rises towards the right.

HIGHER LEVEL

Contents

Topic 1: Number and algebra – HL	2
Topic 2: Functions – HL	3
Topic 3: Geometry and trigonometry – HL	4
Topic 4: Statistics and probability – HL	7
Topic 5: Calculus – HL	9

Topic 1: Number and algebra – HL

1.2	<p>The nth term of an arithmetic sequence</p> <p>The sum of n terms of an arithmetic sequence</p>	$u_n = u_1 + (n - 1)d$ $S_n = \frac{n}{2}(2u_1 + (n - 1)d); S_n = \frac{n}{2}(u_1 + u_n)$
1.3	<p>The nth term of a geometric sequence</p> <p>The sum of n terms of a finite geometric sequence</p>	$u_n = u_1 r^{n-1}$ $S_n = \frac{u_1(r^n - 1)}{r - 1} = \frac{u_1(1 - r^n)}{1 - r}, r \neq 1$
1.8	The sum of an infinite geometric sequence	$S_\infty = \frac{u_1}{1 - r}, r < 1$
1.4	Compound interest	$FV = PV \times \left(1 + \frac{r}{100k}\right)^{kn}$ <p>where FV is the future value, PV is the present value, n is the number of years, k is the number of compounding periods per year, $r\%$ is the nominal annual rate of interest</p>
1.5	Exponents and logarithms	$a^x = b \Leftrightarrow x = \log_a b, \text{ where } a > 0, b > 0, a \neq 1$
1.7	<p>Exponents and logarithms</p> <p>Exponential and logarithmic functions</p>	$\log_a xy = \log_a x + \log_a y$ $\log_a \frac{x}{y} = \log_a x - \log_a y$ $\log_a x^m = m \log_a x$ $\log_a x = \frac{\log_b x}{\log_b a}$ $a^x = e^{x \ln a}; \log_a a^x = x = a^{\log_a x} \text{ where } a, x > 0, a \neq 1$
1.9	Binomial theorem $n \in \mathbb{N}$	$(a + b)^n = a^n + {}^n C_1 a^{n-1} b + \dots + {}^n C_r a^{n-r} b^r + \dots + b^n$ ${}^n C_r = \frac{n!}{r!(n-r)!}$

1.10	Combinations	${}^n C_r = \frac{n!}{r!(n-r)!}$
	Permutations	${}^n P_r = \frac{n!}{(n-r)!}$
	Extension of binomial theorem, $n \in \mathbb{Q}$	$(a+b)^n = a^n \left(1 + n \left(\frac{b}{a} \right) + \frac{n(n-1)}{2!} \left(\frac{b}{a} \right)^2 + \dots \right)$
1.12	Complex numbers	$z = a + bi$
1.13	Modulus-argument (polar) and exponential (Euler) form	$z = r(\cos \theta + i \sin \theta) = r e^{i\theta} = r \operatorname{cis} \theta$
1.14	De Moivre's theorem	$[r(\cos \theta + i \sin \theta)]^n = r^n (\cos n\theta + i \sin n\theta) = r^n e^{in\theta} = r^n \operatorname{cis} n\theta$

Topic 2: Functions – HL

2.1	Equations of a straight line	$y = mx + c; ax + by + d = 0; y - y_1 = m(x - x_1)$
	Gradient formula	$m = \frac{y_2 - y_1}{x_2 - x_1}$
2.6	Axis of symmetry of the graph of a quadratic function	$f(x) = ax^2 + bx + c \Rightarrow$ axis of symmetry is $x = -\frac{b}{2a}$
2.7	Solutions of a quadratic equation	$ax^2 + bx + c = 0 \Rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}, a \neq 0$
	Discriminant	$\Delta = b^2 - 4ac$
2.12	Sum and product of the roots of polynomial equations of the form $\sum_{r=0}^n a_r x^r = 0$	Sum is $-\frac{a_{n-1}}{a_n}$; product is $\frac{(-1)^n a_0}{a_n}$

Topic 3: Geometry and trigonometry – HL

Prior learning – HL

Area of a parallelogram	$A = bh$, where b is the base, h is the height
Area of a triangle	$A = \frac{1}{2}(bh)$, where b is the base, h is the height
Area of a trapezoid	$A = \frac{1}{2}(a + b)h$, where a and b are the parallel sides, h is the height
Area of a circle	$A = \pi r^2$, where r is the radius
Circumference of a circle	$C = 2\pi r$, where r is the radius
Volume of a cuboid	$V = lwh$, where l is the length, w is the width, h is the height
Volume of a cylinder	$V = \pi r^2 h$, where r is the radius, h is the height
Volume of a prism	$V = Ah$, where A is the area of cross-section, h is the height
Area of the curved surface of a cylinder	$A = 2\pi r h$, where r is the radius, h is the height
Distance between two points (x_1, y_1) and (x_2, y_2)	$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$
Coordinates of the midpoint of a line segment with endpoints (x_1, y_1) and (x_2, y_2)	$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

3.1	Distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2)	$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$
	Coordinates of the midpoint of a line segment with endpoints (x_1, y_1, z_1) and (x_2, y_2, z_2)	$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right)$
	Volume of a right-pyramid	$V = \frac{1}{3}Ah$, where A is the area of the base, h is the height

	Volume of a right cone	$V = \frac{1}{3}\pi r^2 h$, where r is the radius, h is the height
	Area of the curved surface of a cone	$A = \pi r l$, where r is the radius, l is the slant height
	Volume of a sphere	$V = \frac{4}{3}\pi r^3$, where r is the radius
	Surface area of a sphere	$A = 4\pi r^2$, where r is the radius
3.2	Sine rule	$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
	Cosine rule	$c^2 = a^2 + b^2 - 2ab \cos C$; $\cos C = \frac{a^2 + b^2 - c^2}{2ab}$
	Area of a triangle	$A = \frac{1}{2}ab \sin C$
3.4	Length of an arc	$l = r\theta$, where r is the radius, θ is the angle measured in radians
	Area of a sector	$A = \frac{1}{2}r^2\theta$, where r is the radius, θ is the angle measured in radians
3.5	Identity for $\tan \theta$	$\tan \theta = \frac{\sin \theta}{\cos \theta}$
3.6	Pythagorean identity	$\cos^2 \theta + \sin^2 \theta = 1$
	Double angle identities	$\sin 2\theta = 2 \sin \theta \cos \theta$ $\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$
3.9	Reciprocal trigonometric identities	$\sec \theta = \frac{1}{\cos \theta}$ $\operatorname{cosec} \theta = \frac{1}{\sin \theta}$
	Pythagorean identities	$1 + \tan^2 \theta = \sec^2 \theta$ $1 + \cot^2 \theta = \operatorname{cosec}^2 \theta$

3.10	Compound angle identities Double angle identity for tan	$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$ $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$ $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$ $\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$
3.12	Magnitude of a vector	$ \mathbf{v} = \sqrt{v_1^2 + v_2^2 + v_3^2}, \text{ where } \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$
3.13	Scalar product Angle between two vectors	$\mathbf{v} \cdot \mathbf{w} = v_1 w_1 + v_2 w_2 + v_3 w_3, \text{ where } \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}, \mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$ $\mathbf{v} \cdot \mathbf{w} = \mathbf{v} \mathbf{w} \cos \theta, \text{ where } \theta \text{ is the angle between } \mathbf{v} \text{ and } \mathbf{w}$ $\cos \theta = \frac{v_1 w_1 + v_2 w_2 + v_3 w_3}{ \mathbf{v} \mathbf{w} }$
3.14	Vector equation of a line Parametric form of the equation of a line Cartesian equations of a line	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$ $x = x_0 + \lambda l, y = y_0 + \lambda m, z = z_0 + \lambda n$ $\frac{x - x_0}{l} = \frac{y - y_0}{m} = \frac{z - z_0}{n}$
3.16	Vector product Area of a parallelogram	$\mathbf{v} \times \mathbf{w} = \begin{pmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - v_1 w_3 \\ v_1 w_2 - v_2 w_1 \end{pmatrix}, \text{ where } \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}, \mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$ $ \mathbf{v} \times \mathbf{w} = \mathbf{v} \mathbf{w} \sin \theta, \text{ where } \theta \text{ is the angle between } \mathbf{v} \text{ and } \mathbf{w}$ $A = \mathbf{v} \times \mathbf{w} \text{ where } \mathbf{v} \text{ and } \mathbf{w} \text{ form two adjacent sides of a parallelogram}$
3.17	Vector equation of a plane Equation of a plane (using the normal vector) Cartesian equation of a plane	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{b} + \mu \mathbf{c}$ $\mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$ $ax + by + cz = d$

Topic 4: Statistics and probability – HL

4.2	Interquartile range	$IQR = Q_3 - Q_1$
4.3	Mean, \bar{x} , of a set of data	$\bar{x} = \frac{\sum_{i=1}^k f_i x_i}{n}$, where $n = \sum_{i=1}^k f_i$
4.5	Probability of an event A	$P(A) = \frac{n(A)}{n(U)}$
	Complementary events	$P(A) + P(A') = 1$
4.6	Combined events	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$
	Mutually exclusive events	$P(A \cup B) = P(A) + P(B)$
	Conditional probability	$P(A B) = \frac{P(A \cap B)}{P(B)}$
	Independent events	$P(A \cap B) = P(A)P(B)$
4.7	Expected value of a discrete random variable X	$E(X) = \sum_{i=1}^k x_i P(X = x_i)$
4.8	Binomial distribution $X \sim B(n, p)$	
	Mean	$E(X) = np$
	Variance	$\text{Var}(X) = np(1-p)$
4.12	Standardized normal variable	$z = \frac{x - \mu}{\sigma}$
4.13	Bayes' theorem	$P(B A) = \frac{P(B)P(A B)}{P(B)P(A B) + P(B')P(A B')}$ $P(B_i A) = \frac{P(B_i)P(A B_i)}{P(B_1)P(A B_1) + P(B_2)P(A B_2) + P(B_3)P(A B_3)}$

<p>4.14</p>	<p>Variance σ^2</p> <p>Standard deviation σ</p> <p>Linear transformation of a single random variable</p> <p>Expected value of a continuous random variable X</p> <p>Variance</p> <p>Variance of a discrete random variable X</p> <p>Variance of a continuous random variable X</p>	$\sigma^2 = \frac{\sum_{i=1}^k f_i (x_i - \mu)^2}{n} = \frac{\sum_{i=1}^k f_i x_i^2}{n} - \mu^2$ $\sigma = \sqrt{\frac{\sum_{i=1}^k f_i (x_i - \mu)^2}{n}}$ $E(aX + b) = aE(X) + b$ $\text{Var}(aX + b) = a^2 \text{Var}(X)$ $E(X) = \mu = \int_{-\infty}^{\infty} x f(x) dx$ $\text{Var}(X) = E[(X - \mu)^2] = E(X^2) - [E(X)]^2$ $\text{Var}(X) = \sum (x - \mu)^2 P(X = x) = \sum x^2 P(X = x) - \mu^2$ $\text{Var}(X) = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx = \int_{-\infty}^{\infty} x^2 f(x) dx - \mu^2$
--------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Topic 5: Calculus – HL

5.12	Derivative of $f(x)$ from first principles	$y = f(x) \Rightarrow \frac{dy}{dx} = f'(x) = \lim_{h \rightarrow 0} \left(\frac{f(x+h) - f(x)}{h} \right)$
5.3	Derivative of x^n	$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$
5.6	Derivative of $\sin x$	$f(x) = \sin x \Rightarrow f'(x) = \cos x$
	Derivative of $\cos x$	$f(x) = \cos x \Rightarrow f'(x) = -\sin x$
	Derivative of e^x	$f(x) = e^x \Rightarrow f'(x) = e^x$
	Derivative of $\ln x$	$f(x) = \ln x \Rightarrow f'(x) = \frac{1}{x}$
	Chain rule	$y = g(u), \text{ where } u = f(x) \Rightarrow \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
	Product rule	$y = uv \Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$
	Quotient rule	$y = \frac{u}{v} \Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
5.15	Standard derivatives	
	$\tan x$	$f(x) = \tan x \Rightarrow f'(x) = \sec^2 x$
	$\sec x$	$f(x) = \sec x \Rightarrow f'(x) = \sec x \tan x$
	$\operatorname{cosec} x$	$f(x) = \operatorname{cosec} x \Rightarrow f'(x) = -\operatorname{cosec} x \cot x$
	$\cot x$	$f(x) = \cot x \Rightarrow f'(x) = -\operatorname{cosec}^2 x$
	a^x	$f(x) = a^x \Rightarrow f'(x) = a^x (\ln a)$
	$\log_a x$	$f(x) = \log_a x \Rightarrow f'(x) = \frac{1}{x \ln a}$
	$\arcsin x$	$f(x) = \arcsin x \Rightarrow f'(x) = \frac{1}{\sqrt{1-x^2}}$
	$\arccos x$	$f(x) = \arccos x \Rightarrow f'(x) = -\frac{1}{\sqrt{1-x^2}}$
	$\arctan x$	$f(x) = \arctan x \Rightarrow f'(x) = \frac{1}{1+x^2}$

5.9	<p>Acceleration</p> <p>Distance travelled from t_1 to t_2</p> <p>Displacement from t_1 to t_2</p>	$a = \frac{dv}{dt} = \frac{d^2s}{dt^2}$ $\text{distance} = \int_{t_1}^{t_2} v(t) dt$ $\text{displacement} = \int_{t_1}^{t_2} v(t) dt$
5.5	<p>Integral of x^n</p> <p>Area between a curve $y = f(x)$ and the x-axis, where $f(x) > 0$</p>	$\int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1$ $A = \int_a^b y dx$
5.10	Standard integrals	$\int \frac{1}{x} dx = \ln x + C$ $\int \sin x dx = -\cos x + C$ $\int \cos x dx = \sin x + C$ $\int e^x dx = e^x + C$
5.15	Standard integrals	$\int a^x dx = \frac{1}{\ln a} a^x + C$ $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C$ $\int \frac{1}{\sqrt{a^2 - x^2}} dx = \arcsin\left(\frac{x}{a}\right) + C, x < a$
5.16	Integration by parts	$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx \text{ or } \int u dv = uv - \int v du$

5.11	Area of region enclosed by a curve and x -axis	$A = \int_a^b y dx$
5.17	Area of region enclosed by a curve and y -axis Volume of revolution about the x or y -axes	$A = \int_a^b x dy$ $V = \int_a^b \pi y^2 dx$ or $V = \int_a^b \pi x^2 dy$
5.18	Euler's method Integrating factor for $y' + P(x)y = Q(x)$	$y_{n+1} = y_n + h \times f(x_n, y_n)$; $x_{n+1} = x_n + h$, where h is a constant (step length) $e^{\int P(x) dx}$
5.19	Maclaurin series Maclaurin series for special functions	$f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \dots$ $e^x = 1 + x + \frac{x^2}{2!} + \dots$ $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$ $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$ $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$ $\arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$